

Trophy Rack Spring/Summer Deer Mix

Foxtail Millet

- Foxtail millet is an annual grass with slim, vertical, leafy stems which can reach a height of 120–200 cm (3.9–6.6 ft).
- The seedhead is a dense, hairy panicle 5–30 cm (2.0–12 in) long.
- The small seeds, around 2 mm (less than 1/8 in.) in diameter, are encased in a thin, papery hull which is easily removed in threshing. Seed color varies greatly between varieties



Red Clover

- It is an herbaceous, short-lived perennial plant, variable in size, growing to 20–80 cm tall.
- The leaves are alternate, trifoliate (with three leaflets), each leaflet 15–30 mm long and 8–15 mm broad, green with a characteristic pale crescent in the outer half of the leaf; the petiole is 1–4 cm long, with two basal stipules.
- The flowers are dark pink with a paler base, 12–15 mm long, produced in a dense inflorescence.



Ladino Clover

- It is a herbaceous, perennial plant. It is low growing, with heads of whitish flowers, often with a tinge of pink or cream that may come on with the aging of the plant.
- The heads are generally 1.5–2 centimeters (0.6–0.8 in) wide, and are at the end of 7-cm (2.8-in) peduncles or flower stalks.
- The leaves, which by themselves form the symbol known as shamrock, are trifoliolate, smooth, elliptic to egg-shaped and long-petioled.
- The stems function as stolons, so white clover often forms mats, with the stems creeping as much as 18 cm (7.1 in) a year, and rooting at the nodes.



Buckwheat

- Buckwheat plants grow quickly, beginning to produce seed in about 6 weeks and ripening at 10 to 11 weeks. They grow 30 to 50 inches (75 to 125 cm) tall.



White Clover

- It is a herbaceous, perennial plant. It is low growing, with heads of whitish flowers, often with a tinge of pink or cream that may come on with the aging of the plant.
- The heads are generally 1.5–2 centimeters (0.6–0.8 in) wide, and are at the end of 7-cm (2.8-in) peduncles or flower stalks.
- The leaves, which by themselves form the symbol known as shamrock, are trifoliolate, smooth, elliptic to egg-shaped and long-petioled.
- The stems function as stolons, so white clover often forms mats, with the stems creeping as much as 18 cm (7.1 in) a year, and rooting at the nodes.



Birdsfoot Trefoil

- Most species have leaves with five leaflets; two of these are at the extreme base of the leaf, with the other three at the tip of a naked midrib.
- Some species have pinnate leaves with up to 15 leaflets. The flowers are in clusters of 3-10 together at the apex of a stem with some basal leafy bracts; they are pea-flower shaped, usually vivid yellow, but occasionally orange or red. The seeds develop in three or four straight, strongly diverging pods, which together make a shape reminiscent of the diverging toes of a small bird, leading to the common name "bird's-foot"



Rape Seed

- A low growing, dense, broadleaf plant.

**Chicory**

- When flowering, chicory has a tough, grooved, and more or less hairy stem, from 30 to 100 centimeters (10 to 40 in) tall.
- The leaves are stalked, lanceolate and unlobed.
- The flower heads are 2 to 4 centimeters (0.79 to 1.6 in) wide, and usually bright blue, rarely white or pink. There are two rows of involucral bracts; the inner are longer and erect, the outer are shorter and spreading. It flowers from July until October.

